

Metabolism:
**Estradiol stimulates glucose metabolism via
6-phosphofructo-2-kinase (PFKFB3)**

METABOLISM

Yoannis Imbert-Fernandez, Brian F. Clem,
Julie O'Neal, Daniel A. Kerr, Robert
Spaulding, Lilibeth Lanceta, Amy L. Clem,
Sucheta Telang and Jason Chesney
J. Biol. Chem. published online February 10, 2014

Access the most updated version of this article at doi: [10.1074/jbc.M113.529990](https://doi.org/10.1074/jbc.M113.529990)

Find articles, minireviews, Reflections and Classics on similar topics on the [JBC Affinity Sites](#).

Alerts:

- [When this article is cited](#)
- [When a correction for this article is posted](#)

[Click here](#) to choose from all of JBC's e-mail alerts

This article cites 0 references, 0 of which can be accessed free at
<http://www.jbc.org/content/early/2014/02/10/jbc.M113.529990.full.html#ref-list-1>

Estradiol Stimulates Glucose Metabolism via 6-Phosphofructo-2-Kinase (PFKFB3)

Yoannis Imbert-Fernandez, Brian F. Clem, Julie O'Neal, Daniel A. Kerr, Robert Spaulding, Lilibeth Lanceta, Amy L. Clem, Sucheta Telang and Jason Chesney

James Graham Brown Cancer Center, Division of Medical Oncology and Hematology, Department of Medicine, University of Louisville, Louisville, KY 40202, USA.

***Running title:** PFKFB3 regulation by estradiol

To whom correspondence should be addressed: Jason Chesney, Department of Medicine, James Graham Brown Cancer Center, University of Louisville, 505 S. Hancock St. Louisville, KY 40202, Tel.: (502) 852-3679; Email: jason.chesney@louisville.edu

Keywords: 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase, PFKFB3, estrogen, estrogen receptor, breast cancer, glycolysis, metabolism.

Background: The regulation of glucose metabolism by estradiol is poorly defined.

Results: We find that estradiol stimulates glucose metabolism in part by stimulating the production of fructose-2,6-bisphosphate by PFKFB3.

Conclusion: PFKFB3 is a downstream target of estradiol required to stimulate glucose metabolism.

Significance: Combined targeting of PFKFB3 and estrogen receptor may prove beneficial to ER+ stage IV breast cancer patients.

Abstract

Estradiol (E2) administered to estrogen receptor-positive (ER+) breast cancer patients stimulates glucose uptake by tumors. Importantly, this E2-induced metabolic flare is predictive of the clinical effectiveness of anti-estrogens and, as a result, downstream metabolic regulators of E2 are expected to have utility as targets for the development of anti-breast cancer agents. The family of 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatases (PFKFB1-4) control glycolytic flux via their product, fructose-2,6-bisphosphate (F26BP), which activates 6-phosphofructo-1-kinase (PFK-1). We postulated that E2 might promote PFKFB3 expression, resulting in

increased F26BP and glucose uptake. We demonstrate that PFKFB3 expression is highest in stage III lymph node metastases relative to normal breast tissues and that exposure of human MCF-7 breast cancer cells to E2 causes a rapid increase in ¹⁴C-glucose uptake and glycolysis that is coincident with an induction of PFKFB3 mRNA (via ER binding to its promoter), protein expression and the intracellular concentration of its product, F26BP. Importantly, selective inhibition of PFKFB3 expression and activity using siRNA or a PFKFB3 inhibitor markedly reduces the E2-mediated increase in F26BP, ¹⁴C-glucose uptake and glycolysis. Furthermore, co-treatment of MCF-7 cells with the PFKFB3 inhibitor and the anti-estrogen, ICI 182,780, synergistically induces apoptotic cell death. These findings demonstrate for the first time that the estrogen receptor directly promotes PFKFB3 mRNA transcription which, in turn, is required for the glucose metabolism and survival of breast cancer cells. Importantly, these results provide essential pre-clinical information that may allow for the ultimate design of combinatorial trials of PFKFB3 antagonists with anti-estrogen therapies in ER+ stage IV breast cancer patients.

Introduction

Increased uptake and metabolism of glucose as a source of energy and biosynthetic precursors is a common feature of primary and metastatic breast adenocarcinomas. As evidence, positron emission tomographic studies with 2- ^{18}F -fluoro-2-deoxy-glucose (FDG-PET) have consistently demonstrated that human breast tumors display a marked increase in glucose uptake *in vivo* relative to adjacent normal breast tissues (1-5). Consequently, FDG-PET scans have become widely used by medical oncologists to assess the extent of disease and clinical responses to chemotherapies in breast cancer patients. Importantly, objective responses to aromatase inhibitors or ER antagonists in ER+ breast cancer patients can be predicted by the magnitude of E2-stimulated 2- ^{18}F -fluoro-2-deoxyglucose (FDG) uptake suggesting that glucose metabolism is an essential clinical target for anti-estrogen therapies (6). Although the stimulation of glucose metabolism by E2 has recently been found to be dependent on the PI3K-AKT pathway (7), relatively little is known about the precise downstream effectors required for E2 to stimulate glucose metabolism in breast cancer cells.

6-Phosphofructo-1-kinase (PFK-1) is the first committed step of glycolysis and a key metabolic sensor of the entire pathway since this enzyme is allosterically inhibited by several products of glucose metabolism including ATP, citrate and H^+ ions. The most potent activator of PFK-1 is a shunt product of glycolysis, fructose-2,6-bisphosphate (F26BP), which can override the allosteric inhibition by ATP and stimulate PFK-1, glucose uptake and flux through the entire glycolytic pathway (8,9). The intracellular concentration of F26BP is maintained by a family of four bifunctional enzymes (PFKFB1-4) with fructose 6-phosphate kinase and F26BP phosphatase activities (10). The PFKFB3 family member is of particular interest since this enzyme: (i) has a very high kinase:bisphosphatase ratio

thus overwhelmingly favoring F26BP synthesis and activation of glycolysis; and (ii) is over-expressed in breast tumors relative to normal mammary epithelial cells *in situ* (10-12).

Although the precise mechanisms for increased PFKFB3 expression in breast cancers are not well understood, PFKFB3 mRNA transcription is promoted by HIF-1 α (13,14) and by the progesterone receptor (PR) (15). Additionally, loss of the tumor suppressor *PTEN* has recently been found to reduce APC/Cdh1-mediated degradation of PFKFB3 (16). The observation that the *PFKFB3* promoter contains putative ER response elements (ERE) (14) prompted us to determine if PFKFB3 is a direct transcriptional target of ER. Considering previous research showing that E2 increases glucose uptake *in vivo* and *in vitro* (6,7), together with the glycolysis-activating function of PFKFB3, we postulated that PFKFB3 may be a downstream effector of E2 that serves to activate the energetic and anabolic pathways required for the survival and growth of breast cancer cells.

Experimental Procedures

Chemicals. β -Estradiol (E2), Cycloheximide (Chx) and Actinomycin D (Act) were purchased from Sigma-Aldrich. ICI 182, 780 (Fulvestrant) was purchased from Tocris. PFK158 was provided by Advanced Cancer Therapeutics. The compound consists of (E)-1-(pyridin-4-yl)-3-(7-(trifluoromethyl)quinolin-2-yl)-prop-2-en-1-one.

Antibodies and Western blot. Antibodies against PFKFB3 and β -actin were purchased from Proteintech and Sigma-Aldrich, respectively. Whole cell lysates were harvested using Pierce Lysis buffer supplemented with protease inhibitors. Protein concentration was determined by the BCA method (Pierce). Proteins were separated on 10 % Mini PROTEAN TGX gels (BioRad) and transferred to PVDF membranes (Millipore). The membranes were blocked with 5 % nonfat milk in TBS-T (0.1 % Tween20) and

immunoblotted with the indicated antibodies. Amersham ECL Western blotting detection reagent (GE Healthcare) was used to detect protein bands. The membranes were visualized on Biomax ML film (Kodak).

Cell culture. MCF-7 and T-47D cells were purchased from American Type Culture Collection (ATCC). MCF-7 cells were maintained in Improved Minimum Essential Medium (IMEM) supplemented with 10% fetal bovine serum (FBS) and Pen-Strep (Invitrogen). T-47D cells were maintained RPMI-1640 Medium supplemented with 10% FBS and 0.2 Units/ml bovine insulin. Cells were serum starved (5% dextran-coated charcoal stripped FBS) in phenol red-free media for 72 hours prior to E2 treatments.

RNA extraction and Real-Time QRT-PCR. Total RNA was extracted using RNeasy spin columns (Qiagen). Reverse transcription was performed by using random primers and the High-Capacity cDNA Archive Kit (Applied Biosystems). QRT-PCR was performed with the ABI PRISM 7500 using TaqMan primers and probes for PFKFB3 and control gene *b-actin* purchased as Assays-on-Demand™ Gene Expression Products (Applied Biosystems). Analysis and fold differences were determined using the comparative C_T method. Unless otherwise indicated, fold change was calculated from the $\Delta\Delta C_T$ values with the formula $2^{-\Delta\Delta C_T}$. Experiments were repeated at least 3 times.

Immunohistochemistry. Five μ m sections of formalin-fixed and paraffin-embedded tumor tissue were mounted on charged glass slides and dried at 58°C for 60 minutes. Slides were first deparaffinized with xylene then epitope retrieval carried out using citrate buffer in a 2100 Retriever (PickCell Laboratories). The sections were blocked with 10% goat serum for 1 hour, then incubated with primary antibody (anti-PFKFB3 1:50, Proteintech) overnight, followed by an HRP-linked goat anti-rabbit secondary antibody (1:300, Pierce Biotechnology). The sections were

developed with 3,3'-diaminobenzidine tetrahydrochloride (DAB, Vector Laboratories) for 2 min and nuclei counterstained with hematoxylin (Sigma-Aldrich) for 2 min. PBS washes were performed between all steps. The slides were dehydrated in graded alcohols (100%, 95%, and 80% ethanol [vol/vol] in H₂O), cleared in xylene and coverslips attached with Permount (Fisher Scientific). Slides were scanned using a ScanScope XT Digital Slide Scanner (Aperio) and data analyzed with the positive pixel count algorithm (ImageScope, Aperio).

Chromatin Immunoprecipitation (ChIP) assay.

ChIP experiments were performed using the simpleChIP kit (Cell Signaling) and following the manufacture's instructions. In brief, 10⁷ MCF-7 cells/IP were treated with vehicle control EtOH or 10 nM E2 for 1 hour. Chromatin was crosslinked by incubating the cells with 1% formaldehyde for 10 min. Immunoprecipitation of the lysed extracts was performed using the following ChIP grade antibodies: anti-ER α (HC-20), anti-ER β (H-150) from Santa Cruz Biotechnology or normal rabbit IgG (Cell Signaling). The primers used for PCR of the PFKFB3 promoter region containing an ERE were: (−2023 to −2198); CGCGTTATTGACAGACAGGA (FWD)/AGCTCACAGGGAGAGCAGAG (REV), and (−2982 to −3308); CTCCTGAGCTCAAGCAATCC (FWD)/GGGTGCTGAGTGGAATCCTA (REV). As a positive control, primers flanking the established ERE in the human pS2 [trefoil factor 1 (*TFF1*)] gene promoter were used (17). Primers to a distal region (−9196 to −9432); AGTTGAATCGCTGCAAAGGT (FWD)/ACCCAGGTGAAATCAACAGC (REV) of the PFKFB3 promoter were used as a negative control.

Measurement of F26BP. MCF-7 cells were treated with EtOH vehicle control, 10 or 100 nM E2 for the indicated times. Cells were washed in PBS and lysed in NaOH/Tris Acetate by heating at 80°C for 5 min. Lysates were neutralized to pH

7.2 by adding ice-cold acetic acid and HEPES. F26BP concentration was determined using the method described by Schaftingen *et. al.* (9).

Glucose uptake. MCF-7 cells treated with EtOH vehicle control, 10 or 100 nM E2 for the indicated time periods and subsequently placed in glucose-free RPMI 1640 for 30 min. ^{14}C -2-deoxyglucose (0.25 $\mu\text{Ci/mL}$; Perkin Elmer) was added for an additional 15 min and cells were then washed thrice with ice-cold RPMI 1640 containing no glucose. Cell lysates were collected in 500 μL of 0.1% SDS, and scintillation counts (counts/min) were measured using 350 μL of lysate. Protein concentration was determined using the BCA assay (Pierce) per manufacturer's instructions and measured on a Powerwave XS plate reader (Biotek). Counts were normalized to protein concentration.

Glycolysis assay. MCF-7 cells growing in six-well plates were incubated in 500 μL complete media containing 1 mCi $[5\text{-}^3\text{H}]\text{glucose}$ for 60 minutes in 5% CO_2 at 37°C . The media was then collected and centrifuged for 5 min at 8000rpm to pellet any cells. To separate the $[^3\text{H}]\text{H}_2\text{O}$ formed via glycolysis from the $[5\text{-}^3\text{H}]\text{glucose}$ added to the media, an evaporation technique in a sealed system was utilized. Briefly, 150 μL of media aliquots were added in open tubes that were placed upright inside scintillation vials containing 1 mL H_2O . The scintillation vials were sealed and the $[^3\text{H}]\text{H}_2\text{O}$ produced by glycolysis through enolase and released to the media was allowed to equilibrate with the H_2O in the outer vial for 48 hours at 37°C . The amounts of $[^3\text{H}]\text{H}_2\text{O}$ that had diffused into the surrounding H_2O was measured on a Tri-Carb 2910 liquid scintillation analyzer (Perkin-Elmer) and compared to $[^3\text{H}]\text{H}_2\text{O}$ and $[5\text{-}^3\text{H}]\text{glucose}$ standards. Protein concentration was determined using the BCA assay (Pierce) per manufacturer's instructions and measured on a Powerwave XS plate reader (Biotek). Counts were normalized to protein concentration.

Transfection and siRNA. Cells were seeded in six-well plates at a density of 1×10^5 cells/well. Cells were transfected with control siRNA or PFKFB3 siRNA (Invitrogen) using Lipofectamine RNAiMAX reagent (Invitrogen) and allowed to incubate for 48 hours prior to E2 treatment.

PFKFB3 adenovirus expression. pAdenoG-HA (Vector) and PFKFB3-HA adenovirus (Human) were purchased from Applied Biological Materials. For infection experiments, MCF-7 cells were seeded in six-well plates at a density of 1×10^5 cells/well and allowed to attach overnight. The cells were then placed in 1 mL of media containing 1×10^6 viral particles and allowed to incubate for 48 hours.

Annexin V/PI staining. MCF-7 cells were seeded in six-well plates at a density of 1×10^5 cells/well and starved for 72 hours prior to treatment. Cells were incubated with 10 nM E2 and DMSO, 100 nM ICI 182,780, 5 μM PFK158 or the combination of ICI 182,780 and PFK158 for 24 hours. Cells were washed with PBS and stained with Annexin V and Propidium Iodide (BD Pharmingen). Fluorescence was measured using a FACSCalibur (BD Biosciences) and analyzed using FloJo (Tree Star). Annexin V-PI- (live) and Annexin V+/PI+ (late apoptotic) cells were quantified by the frequency of fluorescently labeled cells and statistical significance was assessed by the two-sample T-test (independent variable).

Results

High PFKFB3 Protein Expression in Lymph Node Metastases

Previously, PFKFB3 expression was found to be elevated in primary breast adenocarcinomas (12). We confirmed high PFKFB3 expression in stage II breast adenocarcinomas relative to normal breast tissues using immunohistochemical (IHC) analyses (Fig. 1A). We also examined the metastatic breast adenocarcinoma cells in lymph nodes resected

from patients suffering from stage III breast cancer and observed an incremental increase in PFKFB3 protein expression (Fig. 1A). This increase in PFKFB3 expression in stage II and lymph node metastases then was statistically confirmed using digital IHC image analysis (Fig. 1B).

PFKFB3 expression, F26BP, glucose uptake and glycolysis are induced by E2 in MCF-7 cells

Given that E2 increases the glucose uptake of ER+ breast cancer cells *in vitro* and *in vivo* (6,7) and that the *PFKFB3* promoter has putative EREs (14), we postulated that ER and its ligand, E2, might regulate PFKFB3 mRNA expression. Initially, we examined PFKFB3 mRNA expression by real time RT-PCR after exposure to EtOH (vehicle control), 10 or 100 nM E2 for 1, 3, 6 and 24 hours. We observed a marked increase in PFKFB3 mRNA levels after only 1 hour of exposure to either 10 or 100 nM E2 (Fig. 2A). PFKFB3 mRNA returned to near basal levels after 3 and 6 hours of incubation with E2 but was still statistically increased relative to the vehicle control (Fig. 2A). Importantly, co-culture with the anti-estrogen ICI 182,780 (ICI; 100 nM) blocked the induction of PFKFB3 mRNA indicating that the ER is required for the up-regulation of PFKFB3 mRNA expression (Fig. 2A). In addition, pre-treatment of MCF-7 cells with the transcription inhibitor, Actinomycin D (Act), suppressed the increase in PFKFB3 mRNA indicating that new mRNA synthesis is required whereas pre-treatment with the protein synthesis inhibitor, cycloheximide (Chx), had no effect (Fig. 2A). We next examined endogenous PFKFB3 protein expression in response to E2 and found that PFKFB3 protein significantly increased after 3 hours and 6 hours of exposure to 10 or 100 nM E2 but returned to baseline after 24 hours (Fig. 2C and 2D). Similar to PFKFB3 protein, the steady-state concentration of its product, F26BP, significantly increased after 3 hours and 6 hours of exposure to 10 and 100nM E2 (Fig. 2B). To then monitor the effect of E2 on glucose uptake,

MCF-7 cells were placed in phenol red-free media supplemented with 5% dextran-coated charcoal-stripped serum for 72 hours prior to the addition of E2 and 2-[1-¹⁴C]-deoxy-D-glucose (¹⁴C-2DG). We observed a time dependent increase in glucose uptake in MCF-7 cells exposed to 10 and 100 nM E2 for 3-24 hours (Fig. 2E). We next examined glycolysis by measuring the release of ³H₂O from 5-³H-glucose via enolase which is downstream of PFK-1 in the glycolytic pathway. We found that E2 also caused a marked increase in glycolysis by MCF-7 cells that coincided with the observed increase in glucose uptake (Fig. 2F). Given that the F26BP concentration returns to near baseline 6 hours after stimulation by E2, we suspect that the maintenance of increased glucose metabolism observed at 24 hours is mediated by secondary events that regulate glucose transporters and/or glycolytic enzymes. Importantly, we did not observe an increase in the mRNA expression of several key control steps in glycolysis including glut1-4, hexokinase, PFK-1, and pyruvate kinase at 24 hours (*data not shown*). We also assessed PFKFB3 expression after 10 nM E2 exposure in a second ER+ breast cancer cell line (T-47D) and found that PFKFB3 mRNA and protein were increased 3.9-fold (1 hr) and 3.0-fold (3 hrs), respectively (*data not shown*).

ER α is recruited to the *PFKFB3* gene upon E2 stimulation

We next analyzed the 5' *PFKFB3* promoter sequence using the TRANSFAC software (Biobase) in order to identify putative transcription factors binding sites, including the ER. Our analysis revealed four putative half-site ERE sites located around the -2000 and -3000 regions along with several SP1 sites. To investigate whether the E2-induced increase in PFKFB3 mRNA involves recruitment of the ER to the PFKFB3 promoter we performed ChIP assays using primers that could amplify the -2000, -3000 and a distal region of the *PFKFB3* promoter. We used antibodies against ER α and ER β to immunoprecipitate the chromatin

collected from MCF-7 cells treated with 10 nM E2 for 1 hour and then performed real-time quantitative PCR to determine recruitment to the *PFKFB3* promoter. Our results demonstrate a four-fold enrichment in the recruitment of the ER α to the -2000 region of the *PFKFB3* promoter after 1 hour of E2 exposure (Fig. 3A [real-time PCR] and Fig. 3B [2% agarose gel as a representative example]). We did not observe a significant recruitment of the ER α to the -3000 region or a distal region of the *PFKFB3* gene which is devoid of putative EREs. We also observed no recruitment of the ER β to the *PFKFB3* gene and found that primers recognizing the pS2 promoter, a previously described ER α target (18), amplified the pS2 promoter sequence as a positive control. In summary, these studies demonstrate that stimulation of MCF-7 cells with 10 nM E2 for 1 hour causes an increase in ER α recruitment to the *PFKFB3* promoter.

E2 increases glucose uptake and glycolysis via PFKFB3 in MCF-7 cells

To determine if the stimulation of glucose uptake by E2 requires PFKFB3 expression, we next analyzed E2-stimulated glucose uptake by MCF-7 cells transfected with a control or PFKFB3-specific siRNA. Briefly, MCF-7 cells were transiently transfected with either control siRNA or PFKFB3 siRNA and, 48 hours later, the cells were exposed to vehicle or the physiologically relevant 10 nM E2 for 3 hours followed by incubation with ^{14}C -2DG. We confirmed that PFKFB3 protein expression was decreased by transfection with PFKFB3 siRNA relative to the control siRNA and found that the PFKFB3 siRNA completely blocked the E2-mediated upregulation of PFKFB3 expression (Fig. 4A and 4B). PFKFB3 has the highest kinase:bisphosphatase ratio of the PFKFB family and we postulated that inhibition of PFKFB3 expression would suppress the intracellular F26BP concentration. We found that transfection with the PFKFB3 siRNA (which does not affect the other family members) markedly reduced the F26BP concentration in vehicle and E2-stimulated

MCF-7 cells after 3 hours (Fig. 4C). In order to test the hypothesis that PFKFB3/F26BP is required for the stimulation of glucose uptake and glycolysis by E2, we examined the effect of PFKFB3 siRNA on the ^{14}C -2DG uptake and $^3\text{H}_2\text{O}$ release (*i.e.* glycolysis) stimulated by E2. We found that transfection with the PFKFB3 siRNA reduced basal glucose uptake and glycolysis and completely blocked the stimulation of glucose uptake and glycolysis caused by E2 at 3 hours (Fig. 4D and 4E). Taken together, these data indicate that E2 rapidly stimulates glucose uptake in part by increasing PFKFB3 expression and F26BP.

A small molecule inhibitor of PFKFB3 blocks the E2-mediated increase in glucose uptake and glycolysis in MCF-7 cells

We next postulated that inhibition of the kinase activity of PFKFB3 by a clinical-grade PFKFB3 inhibitor, PFK158, would, like the PFKFB3 siRNA, attenuate the E2-mediated stimulation of F26BP and glucose uptake and glycolysis. Initially, we identified a concentration of PFK158 that does not affect viability after 3 hours of exposure (3 μM ; *data not shown*). We then examined the effect of PFK158 on PFKFB3 protein expression in the absence and presence of 10 nM E2 and observed relatively little effect on the E2-induced expression of PFKFB3 (Fig. 5A and 5B). However, we did observe a marked decrease in the E2-stimulated F26BP (Fig. 5C), ^{14}C -deoxy-glucose uptake (Fig. 5D) and glycolysis (Fig. 5E). These data indicate that selective inhibition of the substrate-binding domain of PFKFB3 is sufficient to completely block the ability of E2 to stimulate glucose uptake. Of note, whereas transfection of unstimulated MCF-7 cells with the PFKFB3 siRNA significantly reduced glucose uptake and glycolysis (Figure 4D, E), exposure of unstimulated MCF-7 cells to PFK158 did not reduce glucose uptake or glycolysis (Fig. 5D, 5E). We believe that this disparity may be due to the relative potency of PFK158 which competitively but not completely inhibits PFKFB3 (unlike the

PFKFB3 siRNA which essentially depletes all PFKFB3 protein).

Forced expression of PFKFB3 increases F26BP, glucose uptake and glycolysis in MCF-7 cells

Given that both PFKFB3 siRNA and PFK158 can suppress the E2-stimulated increase in F26BP, glucose uptake and glycolysis, we sought to determine if PFKFB3 expression by itself would result in similar biochemical changes. We transiently expressed PFKFB3 in MCF-7 cells and found that forced expression of PFKFB3 mRNA (Figure 6A) and protein (Figure 6C and 6D) can increase the steady-state concentration of F26BP (Figure 6B), glucose uptake (Figure 6E) and glycolysis (Figure 6F) in MCF-7 cells.

Effect of ICI 182,780 on PFK158-induced apoptotic death

In order to determine the utility of combining anti-estrogens with PFKFB3 inhibitors, we next examined the effects of 100 nM ICI 182,780 (Fulvestrant), a commonly prescribed pure anti-estrogen compound, a pro-apoptotic concentration of PFK158 (5 μ M at 24 hours) or combined 100 nM ICI 182,780 + 5 μ M PFK158 on apoptosis using PI/Annexin V staining by flow cytometry. We observed that the addition of ICI 182,780 caused a synergistic increase in late apoptotic MCF-7 cells relative to that caused by PFK158 alone after 24 hours of exposure (Fig. 7A and 7B). These data suggest that the addition of PFK158 to anti-estrogen therapeutic agents may increase response rates and perhaps the durability of responses in pre-clinical and clinical studies.

Discussion

In this study, we describe PFKFB3 as an early molecular target of estradiol via direct binding of the ER to the *PFKFB3* promoter. We also show that E2 up-regulates PFKFB3 expression in ER-responsive MCF-7 cells,

increasing the steady-state concentration of F26BP, glucose uptake and glycolysis. Although our study does not characterize all of the potential mechanisms by which E2 increases glucose metabolism, our results do demonstrate that PFKFB3 expression and activity are required to mediate E2-mediated glucose uptake and glycolysis. We postulate that the increase in F26BP by E2 leads to the stimulation of PFK-1 which, in turn, may cause a reduction in the PFK-1 substrate F6P, which is in equilibrium with G6P, an allosteric inhibitor of hexokinase, which itself is required for glucose uptake (19-21). Importantly, the ability of PFKFB3 to regulate glucose uptake has been previously demonstrated by work showing that genetic ablation of the *Pfkfb3* gene reduces F26BP and glucose uptake in transformed cells *in vivo* and *in vitro* (22).

The therapeutic implications of our observation that E2 requires PFKFB3 to stimulate glucose metabolism *in vitro* are two-fold: (i) F26BP may be required for the metabolic flare induced by E2 in breast cancer patients; and (ii) combined inhibition of PFKFB3 and ER functions may be an effective therapeutic strategy for ER+ stage IV breast cancer patients.

The activation of the ER by estrogens serves a critical role in cancer initiation and progression, and their antagonists have proven efficacy in the treatment of breast cancers. Anti-estrogen therapies include selective ER modulators such as tamoxifen that act as partial receptor agonists, estrogen synthesis inhibitors such as aromatase inhibitors, and receptor expression inhibitors such as the anti-estrogen ICI 182,780 (Fulvestrant). Each of these modalities is effective in 50-70% of breast cancer patients initially, but recurrent ER+ breast tumors respond to such anti-estrogen treatments in only ~30% of patients, and durable responses are even rarer (23). Consequently, there is an important need to identify novel ER signaling targets, such as PFKFB3, in order to guide the development of

therapeutic strategies to maximize endocrine responses.

Our observation that E2 increases PFKFB3 expression and F26BP synthesis is the first report that ER signaling regulates PFKFB3 expression and activity. PFKFB3 is a key regulator of glucose metabolism and has proven to be an important target for antineoplastic therapy (24). Additionally, PFKFB3 ChIP experiments demonstrate interaction of the ER α at the half-site EREs located on the -2000 region of the PFKFB3 promoter. Based on these results, we envision the following series of events mediated by exposure of ER+ breast cancer cells to E2: (i) E2 increases PFKFB3 transcription by a mechanism involving ER α occupancy of the PFKFB3 promoter; (ii) the increase in PFKFB3 mRNA translates into an increase in protein and more importantly, an increase in the product, F26BP; and (iii) F26BP activates PFK-1, an essential control enzyme for the entire glycolytic pathway, which reduces F6P and G6P, resulting in less allosteric inhibition of hexokinase and increased glucose uptake (Figure 8).

Conclusions

References

1. Constantinidou, A., Martin, A., Sharma, B., and Johnston, S. R. (2010) Positron emission tomography/computed tomography in the management of recurrent/metastatic breast cancer: a large retrospective study from the Royal Marsden Hospital. *Ann Oncol* **22**, 307-14
2. Martoni, A. A., Zamagni, C., Quercia, S., Rosati, M., Cacciari, N., Bernardi, A., Musto, A., Fanti, S., Santini, D., and Taffurelli, M. (2010) Early (18)F-2-fluoro-2-deoxy-d-glucose positron emission tomography may identify a subset of patients with estrogen receptor-positive breast cancer who will not respond optimally to preoperative chemotherapy. *Cancer* **116**, 805-813
3. Niikura, N., and Ueno, N. T. (2010) The Role of F-FDG-Positron Emission Tomography/Computed Tomography in Staging Primary Breast Cancer. *J Cancer* **1**, 51-53
4. Pennant, M., Takwoingi, Y., Pennant, L., Davenport, C., Fry-Smith, A., Eisinga, A., Andronis, L., Arvanitis, T., Deeks, J., and Hyde, C. (2010) A systematic review of positron emission tomography (PET) and positron emission tomography/computed tomography (PET/CT) for the diagnosis of breast cancer recurrence. *Health Technol Assess* **14**, 1-103

In this study we have found that PFKFB3 and its product, F26BP, are induced by E2 and required for the E2-mediated stimulation of glucose uptake and glycolysis. We also have found that a PFKFB3 inhibitor undergoing phase I trial testing in solid tumor patients, PFK158, causes a synergistic increase in apoptosis by ER+ breast cancer cells when combined with the widely used anti-estrogen ICI 182,780 (Fulvestrant). These observations support the conduct of Investigational New Drug (IND)-enabling pre-clinical toxicity and efficacy studies of combinations of PFK158 and ICI 182,780 in order to facilitate the eventual development of phase I/II trials in ER+ stage IV breast cancer patients.

Acknowledgements

We thank the following funding agencies for their support of these studies: YIF and JO: CDMRP Post-Doctoral Fellowships; ST, NCI 1R01CA140991 and ACS RSG-10-021-01CNE; and JC, NCI 1R01CA149438. We also thank Drs. John Eaton and Otto Grubaw for assisting with the interpretation of our data.

5. Specht, J. M., Kurland, B. F., Montgomery, S. K., Dunnwald, L. K., Doot, R. K., Gralow, J. R., Ellis, G. K., Linden, H. M., Livingston, R. B., Allison, K. H., Schubert, E. K., and Mankoff, D. A. (2010) Tumor metabolism and blood flow as assessed by positron emission tomography varies by tumor subtype in locally advanced breast cancer. *Clin Cancer Res* **16**, 2803-2810
6. Dehdashti, F., Mortimer, J. E., Trinkaus, K., Naughton, M. J., Ellis, M., Katzenellenbogen, J. A., Welch, M. J., and Siegel, B. A. (2009) PET-based estradiol challenge as a predictive biomarker of response to endocrine therapy in women with estrogen-receptor-positive breast cancer. *Breast Cancer Res Treat* **113**, 509-517
7. Ko, B. H., Paik, J. Y., Jung, K. H., and Lee, K. H. (2010) 17beta-estradiol augments 18F-FDG uptake and glycolysis of T47D breast cancer cells via membrane-initiated rapid PI3K-Akt activation. *J Nucl Med* **51**, 1740-1747
8. Van Schaftingen, E., Hue, L., and Hers, H. G. (1980) Fructose 2,6-bisphosphate, the probably structure of the glucose- and glucagon-sensitive stimulator of phosphofructokinase. *The Biochemical journal* **192**, 897-901
9. Van Schaftingen, E., Hue, L., and Hers, H. G. (1980) Control of the fructose-6-phosphate/fructose 1,6-bisphosphate cycle in isolated hepatocytes by glucose and glucagon. Role of a low-molecular-weight stimulator of phosphofructokinase. *The Biochemical journal* **192**, 887-895
10. Yalcin, A., Telang, S., Clem, B., and Chesney, J. (2009) Regulation of glucose metabolism by 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatases in cancer. *Experimental and molecular pathology* **86**, 174-179
11. Sakakibara, R., Kato, M., Okamura, N., Nakagawa, T., Komada, Y., Tominaga, N., Shimojo, M., and Fukasawa, M. (1997) Characterization of a human placental fructose-6-phosphate, 2-kinase/fructose-2,6-bisphosphatase. *J Biochem (Tokyo)* **122**, 122-128
12. Atsumi, T., Chesney, J., Metz, C., Leng, L., Donnelly, S., Makita, Z., Mitchell, R., and Bucala, R. (2002) High expression of inducible 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase (iPFK-2; PFKFB3) in human cancers. *Cancer Res* **62**, 5881-5887
13. Minchenko, A., Leshchinsky, I., Opentanova, I., Sang, N., Srinivas, V., Armstead, V., and Caro, J. (2002) Hypoxia-inducible factor-1-mediated expression of the 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase-3 (PFKFB3) gene. Its possible role in the Warburg effect. *The Journal of biological chemistry* **277**, 6183-6187
14. Obach, M., Navarro-Sabate, A., Caro, J., Kong, X., Duran, J., Gomez, M., Perales, J. C., Ventura, F., Rosa, J. L., and Bartrons, R. (2004) 6-Phosphofructo-2-kinase (pfkfb3) gene promoter contains hypoxia-inducible factor-1 binding sites necessary for transactivation in response to hypoxia. *The Journal of biological chemistry* **279**, 53562-53570
15. Novellasdemunt, L., Obach, M., Millan-Arino, L., Manzano, A., Ventura, F., Rosa, J. L., Jordan, A., Navarro-Sabate, A., and Bartrons, R. (2012) Progestins activate 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase 3 (PFKFB3) in breast cancer cells. *The Biochemical journal* **442**, 345-356
16. Garcia-Cao, I., Song, M. S., Hobbs, R. M., Laurent, G., Giorgi, C., de Boer, V. C., Anastasiou, D., Ito, K., Sasaki, A. T., Rameh, L., Carracedo, A., Vander Heiden, M. G., Cantley, L. C., Pinton, P., Haigis, M. C., and Pandolfi, P. P. (2012) Systemic elevation of PTEN induces a tumor-suppressive metabolic state. *Cell* **149**, 49-62
17. Bourdeau, V., Deschenes, J., Metivier, R., Nagai, Y., Nguyen, D., Bretschneider, N., Gannon, F., White, J. H., and Mader, S. (2004) Genome-wide identification of high-affinity estrogen response elements in human and mouse. *Mol Endocrinol* **18**, 1411-1427

18. Liu, X. F., and Bagchi, M. K. (2004) Recruitment of distinct chromatin-modifying complexes by tamoxifen-complexed estrogen receptor at natural target gene promoters in vivo. *The Journal of biological chemistry* **279**, 15050-15058
19. Gregoriou, M., Cornish-Bowden, A., and Trayer, I. P. (1981) Isotope-exchange evidence for allosteric regulation of hexokinase II by glucose 6-phosphate and for an obligatory addition of substrates. *Biochemical Society transactions* **9**, 62-63
20. Gregoriou, M., Trayer, I. P., and Cornish-Bowden, A. (1983) Isotope-exchange evidence that glucose 6-phosphate inhibits rat-muscle hexokinase II at an allosteric site. *European journal of biochemistry / FEBS* **134**, 283-288
21. Lazo, P. A., and Bosca, L. (1982) Mitochondrial membrane-bound hexokinase of ascites tumor cells. Functional implications of lysine residues studied by modification with imidoesters. *Hoppe Seylers Z Physiol Chem* **363**, 635-641
22. Telang, S., Yalcin, A., Clem, A. L., Bucala, R., Lane, A. N., Eaton, J. W., and Chesney, J. (2006) Ras transformation requires metabolic control by 6-phosphofructo-2-kinase. *Oncogene* **25**, 7225-7234
23. Ingle, J. N. (2013) Pharmacogenomics of endocrine therapy in breast cancer. *Journal of human genetics* **58**, 306-12
24. Clem, B. F., O'Neal, J., Tapolsky, G., Clem, A. L., Imbert-Fernandez, Y., Kerr, D. A., 2nd, Klarer, A., Redman, R., Miller, D. M., Trent, J. O., Telang, S., and Chesney, J. (2013) Targeting 6-Phosphofructo-2-Kinase (PFKFB3) as a Therapeutic Strategy against Cancer. *Molecular cancer therapeutics* **12**, 1461-70

Abbreviations

E2- estradiol; ER- estrogen receptor; F26BP- fructose-2,6-bisphosphate; PFKFB- 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase; ChIP- chromatin immunoprecipitation; PET- positron emission tomography; FDG- 2-[¹⁸F]-fluoro-2-deoxyglucose; PFK-1- 6-Phosphofructo-1-kinase; HIF-1α- hypoxia inducible factor 1 alpha; PTEN- Phosphatase and tensin homolog; ERE- estrogen receptor response element; ¹⁴C-2DG- 2-[1-¹⁴C]-deoxy-D-glucose; F6P- fructose-6-phosphate; G6P- glucose-6-phosphate

Figure Legends

Figure 1. High Expression of PFKFB3 by Metastatic Breast Adenocarcinoma Cells in Lymph Nodes. Four normal breast tissues, stage II primary breast adenocarcinomas and stage III lymph node metastases were analyzed for PFKFB3 expression using IHC (A). Digital IHC image analysis was performed and the ratio of positive pixels to total pixels was quantified for each of the tissue specimens (B). *p* values comparing normal mammary tissues to stage II and stage III lymph node metastases and between stage II and III lymph node metastases are included in the figure.

Figure 2. Regulation of PFKFB3, F26BP, Glucose Uptake and Glycolysis by E2 in MCF-7 cells. MCF-7 cells were stimulated with 10 nM and 100 nM E2 for the indicated times and analyzed for PFKFB3 mRNA by real-time PCR (A), PFKFB3 protein expression by Western blot analysis and densitometry (C and D), F26BP using an enzyme-coupled reaction dependent on NADH oxidation (B),

^{14}C -2DG uptake (E) and glycolysis as measured by the release of $^3\text{H}_2\text{O}$ by enolase (F). Values are the mean $\pm$ SEM of 3 experiments and * indicate p value < 0.01 .

Figure 3. E2 Causes ER α Recruitment to the *PFKFB3* Promoter. ChIP assay experiments were performed with MCF-7 cells treated with EtOH or 10 nM E2 for 1 hour using antibodies to ER α (HC20) or ER β (H150) or with rabbit preimmune serum (IgG) and primers flanking the -2000, -3000 and distal regions (lacking ERE sites) of the *PFKFB3* promoter. Real-time PCR fold results were normalized to input and shown as fold enrichment over IgG IP in EtOH-treated cells from 3 independent ChIP experiments (A). Representative 2% agarose gel using similar conditions and primers is also shown (B). * p value < 0.01 compared to vehicle control.

Figure 4. *PFKFB3* Is Required for E2-Induced Glucose Uptake and Glycolysis. MCF-7 cells were transfected with *PFKFB3* siRNA and allowed to incubate for 48 hours prior to stimulation with 10 nM E2. Shown are *PFKFB3* protein expression by Western blot analysis after 3 hours stimulation with E2 (A), densitometry analysis of relative *PFKFB3*/ β -actin (B), F26BP concentration (C), ^{14}C -DG uptake (D), and glycolysis as measured by the release of $^3\text{H}_2\text{O}$ by enolase (E). Values represent three independent experiments. * p value < 0.01 relative to control siRNA.

Figure 5. *PFK158* Inhibits E2-Induced Glucose Uptake and Glycolysis. MCF-7 cells were simultaneously stimulated with E2 and exposed to 3 μM *PFK158* for 3 hours and then analyzed for *PFKFB3* protein expression (A), densitometry analysis of relative *PFKFB3*/ β -actin (B), F26BP (C), ^{14}C -DG uptake (D) and glycolysis as measured by the release of $^3\text{H}_2\text{O}$ by enolase (E). Values represent three independent experiments. * p value < 0.001 relative to vehicle.




Figure 6. Forced Expression of *PFKFB3* Increases Glucose Uptake and Glycolysis. MCF-7 cells were transduced with adenovirus-encoded *PFKFB3* and then analyzed for *PFKFB3* mRNA (A), F26BP (B), *PFKFB3* protein expression (C), densitometry analysis of relative *PFKFB3*/ β -actin (D), ^{14}C -DG uptake (E) and glycolysis as measured by the release of $^3\text{H}_2\text{O}$ by enolase (F). Values represent three independent experiments. * p value < 0.001 relative to vehicle.

Figure 7. Synergistic Pro-Apoptotic Effect of Combined Treatment with ICI 182,780/*PFK158*. MCF-7 cells were exposed to 100 nM ICI 182,780, 5 μM *PFK158* or both agents for 24 hours and analyzed for PI/Annexin V staining by flow cytometry (A). Absolute cell numbers for double negative and double positive cells were quantified (B).

Figure 8. Model of the Regulation of *PFKFB3* by E2. In this schematic, we summarize our model whereby E2 binds and activates ER, translocates to the nucleus and promotes *PFKFB3* mRNA transcription. Translated *PFKFB3* produces F26BP which allosterically activates PFK-1 and stimulates glucose uptake and the entire glycolytic pathway.

Figure 1

A

Normal
BreastStage II
Breast
AdenoCaStage III
Lymph Node
Metastases

B





Figure 2






A**B**

Figure 4






Figure 6









Figure 7





Figure 8

